

The Preliminary Formulation of *Zea Mays* as Exercise Recovery Gel

Azimah Ahmad¹, Khairunizah Hazila Khalid²

¹Faculty of Medicine and Defense Health, National Defence University of Malaysia, Kem Sungai Besi, 57000 Kuala Lumpur, Malaysia

²Food Technology Research Center, Malaysian Agriculture Research & Development Institute (MARDI), 43400 Serdang, Selangor, Malaysia

Corresponding authors: azimah@upnm.edu.my

ABSTRACT

Recovery nutrition play an important role in body adaptation towards exercise and sports performance. Combination of different saccharides and amino acid content fasten recovery result which promotes exercise adaptation and prepared athletes' body for next session of exercise. There are abundant of recovery supplement available but most of it was milk base products which could contribute to lactose intolerance negative effect among consumers. Therefore, *Zea mays* (*ZM*) which is plant based is propose as an alternative food supplement for exercise recovery choices. *ZM* kernel was grind for the juice and homogenized with sugar, fructose, xanthan gum and pectin. The mixture been cook for 20 minutes using double boiler at 95°C. The gel sample then fill in the gel pack and keep at room temperature for shelve live assessment. Proximate analysis showed the presence of sucrose (20.5g/100g), fructose (9.2g/100g) and total carbohydrate is 51.5g/100g. Essential amino acids that presence is arginine (1.75%w/w), histidine (0.52%w/w), isoleucine (0.17%w/w), lysine (0.08%w/w), threonine (0.15%w/w) and valine (0.31%w/w). After 30 days at room temperature, the *ZM* gel showed same texture, colour, smell and taste. Therefore the preliminary formulation of *ZM* gel showed a possible innovation for local exercise recovery gel.

Keywords: *recovery, zea mays, sweet corn, exercise, supplement*

INTRODUCTION

Exercise and physical activity will result in fluid loss, glycogen depleted and muscle soreness. Therefore, structured recovery plan with appropriate nutrient content could promote recovery process thus increase body muscle adaptation following exercise. Nutritional strategies post exercise could promote muscle glycogen restoration, fluid loss replacement, synthesize new proteins to counteract both catabolic state and exercise-induced damage, and improve immune system response (Burke, 2010, Zoorob et al, 2013, Ivy & Ferguson-Stegall, 2013). Studies showed that muscle glycogen could be restored within 24 hours when adequate amount of carbohydrate are ingested (Beelen et al., 2010), while

recovery process shows better result when protein is ingested together (Burke et al., 2004), thus helps in alleviating muscle soreness (McBrier et al. 2010). Review by Bonilla et al, 2020 summary that using 4R's approached to optimized exercise recovery. 4R's referring to rehydration, refuel, repair and rest which the recommended for each athletes applied it every training and competition session for better adaptation and recovery process. In order to meet the rehydration, refuel and repair, customized nutrient of food product are needed to ease achieve the recovery objective. Therefore, nutrient dense food or product that contain recovery nutrients are needed.

There is abundant food product or supplement for recovery are available but most of it contain milk base as to meet the recovery requirement. Unfortunately, there are many people unable to digest milk due to lactose intolerance. A study done by Asmawi and colleagues (2006) have shown that the prevalence of lactose intolerance was high in Asian population with 88% in Malays, 91% in Chinese, and 83% in Indians. Similarly, more than half of the studied population in Thai (Densupsoonton et al., 2004) and Singapore (Bolin et al. 1970) suffer from this digestive problem. Pelly & Burkart (2013) conducted a survey on dietary regimens among athletes competing in 2010 Commonwealth Games in Delhi reported that 10% of the athletes from Southeast Asia countries were likely to follow a low and/or lactose-free diet to avoid the symptoms of lactose intolerance. Based on this statistic, it is imperative for manufacturers and researchers to produce lactose-free recovery supplement. Therefore this study is to purpose Zea mays or sweet corn as a plant based alternative to innovate the recovery gel as recovery choices that could meet the recovery need as well as lactose-free product.

Materials and Methods

Zea mays L. (ZM) or sweet corn (Nelson's Franchise Sdn. Bhd) was N28 specie that produces succulent kernels with a high sweetness level been used to formulate the recovery gel. Corn kernel was brought in frozen form and keep in low temperature in the ice box during travelling before keeping it at the MARDI lab freezer. *ZM* kernel was defrost by rinse the kernel under running water. *ZM* kernel was grind for the juice and homogenized with sugar, fructose, xanthan gum and pectin. The mixture been cook for 20 minutes using double boiler at 95°C. The gel sample then fill in the gel pack and keep at room temperature for shelve live assessment. Some of the sample was sent to lab for proximate analysis.




Figure 1: Recovery gel flow chart preparation

Result and Discussion

Proximate analysis was done to determine the nutritional profile of *ZM* recovery gel. Result showed that formulated recovery gel contains 51.5g of carbohydrate, 1.4g of protein and 1.0g of fat with total calorie of 221kcal. Table 1 showed the list of proximate analysis of the *ZM* recovery gel.

Table 1. Proximate analysis of *Zea mays* gel

Nutrient	Value (per 100g)
Calorie	221
Total Carbohydrate	51.5
Protein	1.4
Fat	1.0
Ash	0.4
Moister	45.7

Zea mays gel contain high energy and the protein present gave a positive requirement to fulfil recovery need. As it has an original sweetness level, the additional sugar is the purpose for manufacturing. Gel form is chosen to ease the intake of *Zea mays* during or after exercise with higher content of nutrient value. The process of producing *Zea mays* gel showed promising local product that could be used as a choice for exercise recovery purposes.

Sample then was analysed for type of saccharides and amino acid present as both of this nutrient have a role to fasten recovery process. The list of saccharides and amino acid content as presented in Table 2.

Table 2. Saccharides content in *Zea mays* gel

Nutrient	Value (per 100g)
Fructose	9.2
Lactose	ND<0.1
Sucrose	20.5
Maltose	ND<0.1
Glucose	ND<0.1

The present of fructose and sucrose showed a good combination of saccharides. Different saccharides used different pathway to produce energy. Study by Jentjens et al, 2014 showed that when fructose and glucose are ingested simultaneously at high rates during cycling exercise, exogenous carbohydrate oxidation rates can reach peak values of ~1.3

g/min compared to when glucose alone only at 0.8g/min. the result support the *Zea mays* saccharide content in promoting sugar uptake in the cell.

Essential amino acids that presence is arginine (1.75%w/w), histidine (0.52%w/w), lysine (0.08%w/w), threonine (0.15%w/w) and valine (0.31%w/w). While branched-chain amino acid (BCAA) such as leucine, isoleucine, and valine also present in minimum amount. BCAA can be oxidized in skeletal muscle when sufficient protein is available whereby the present gel contains moderate protein. It has been discovered that BCAA have beneficial effects in reducing exercise-induced muscle damage effects and promoting muscle–protein synthesis (Negro et al., 2008). In most of commercially available BCAA supplements, at least 5 g of BCAA is needed to produce the beneficial effects of BCAA supplementation. Meanwhile, the suggested ratio for leucine:isoleucine:valine is 2:1:1 or 3:1:1 (Kephart et al., 2016) or leucine:valine:isoleucine is 2:1:1 (Bongiovanni et al, 2020). Similar ratio of the BCAA amino acids was obtained in this study which was 2:1:1. Therefore, this finding showed that our product has the potential to be developed as an alternative source of BCAA for muscle recovery purpose.

Protein had been recommended to enhance recovery rate as well as promoting exercise adaptation. The combination of carbohydrate and protein could result in synergistic effect on muscle recovery (Belen at al, 2010). The present of amino acid in *Zea mays* which is plant sources could give an alternative for those who practicing vegan or looking for milk alternative product. As lactose intolerance prevalence are high, the innovation of plant-based product that have similar of nearly the contain of milk sources-which had proven to promote recovery, could gave a positive sight to promote this product (Jusoh, Ahmad & Tengah, 2019).

Table 3: Amino acid content in *Zea mays* gel

Nutrient	Value (%w/w)
Arginine	1.79
Aspartic Acid	0.18
Alanine	0.08
Glutamic Acid	0.33
Glycine	0.27
Histidine	0.52
Isoleucine	0.17
Leucine	0.19
Lysine	0.08
Methionine	ND<0.01
Phenylalanine	ND<0.01

Proline	0.47
Serine	0.14
Tyrosine	0.23
Threonine	0.15
Valine	0.13

CONCLUSION

Zea mays gel nutritional content showed promising option as an alternative recovery gel. This plant source product could be a great choices for those with lactose intolerance issue without compromising the recovery need. At the same time sweet corn used are from local plantation that could boost the agricultural activity as well with the new product innovation.

ACKNOWLEDGEMENT

We would like to thank our sponsor, Universiti Pertahanan Nasional Malaysia for the financial assistance that makes this study possible (UPNM/2020/GPJP/SP/2). Not to forget MARDI staffs that help in finding the best final form of gel formulation and the collaboration to use their equipment in order to achieve this study objective.

REFERENCES

- Asmawi, M. Z., Seppo, L., Vapaatalo, H., Korpela, R. 2006. Hypolactasia & lactase intolerance among three ethnic groups in Malaysia. *Indian J. Med. Res.* 124(6), 697-704
- Beelen, M., Burke, L. M., Gibala, M. J., Van Loon, J. C. 2010. Nutritional strategies to promote post-exercise recovery. *Int. J. Sport Nutri and Exer. Metab.* 20, 515-532.
- Bongiovanni, T., Genovesi, F., Nemmer, M., Carling, C., Alberti, G., & Howatson, G. (2020). Nutritional interventions for reducing the signs and symptoms of exercise-induced muscle damage and accelerate recovery in athletes: current knowledge, practical application and future perspectives. *European Journal of Applied Physiology*, 120, 1965-1996.
- Bonilla, D. A., Pérez-Idárraga, A., Odriozola-Martínez, A., & Kreider, R. B. (2020). The 4R's Framework of Nutritional Strategies for Post-Exercise Recovery: A Review with Emphasis on New Generation of Carbohydrates. *International Journal of Environmental Research and Public Health*, 18(1), 103.
- Burke, L. *Nutricion en el Deporte/Nutrition in Sport: Un Enfoque Práctico/a Practical Approach*; Editorial Médica Panamericana S.A.: Madrid, Spain, 2010.
- Densupsoontom, N., Jiranpiyo, P., Thamonsiri, N., Wongam, R. 2004. Lactose intolerance in Thai adults. *J. Med. Assoc. Thai.* 87(12), 1501-1505.
- Ivy, J.L.; Ferguson-Stegall, L.M. Nutrient Timing. *Am. J. Lifestyle Med.* **2013**, *8*, 246–259.
- Jusoh, N., Ahmad, A., & Tengah, R. Y. (2019). Evaluation of nutritive values and consumer acceptance of sweet corn (*Zea mays*) juice as a recovery beverage for exercising people. *Malays. J. Fundam. Appl. Sci.*, 15, 504-507.

- Negro, M., Giardina, S., Marzani, B., Marzatico, F. 2008. Branch-chain amino acid supplementation does not enhance athletic performance but affects muscle recovery and the immune system. *J. Sports Med. Phys. Fitness*. 48 (3), 347-351
- Zoorob, R.; Parrish, M.-E.E.; O'Hara, H.; Kalliny, M. Sports Nutrition Needs. *Prim. Care* **2013**, *40*, 475–486.